


Spatial population genetics in heavily managed species: Separating patterns of historical translocation from contemporary gene flow in white-tailed deer

Tyler K. Chafin¹  | Zachery D. Zbinden¹ | Marlis R. Douglas¹ | Bradley T. Martin¹ | Christopher R. Middaugh² | M. Cory Gray² | Jennifer R. Ballard² | Michael E. Douglas¹

¹Department of Biological Sciences, University of Arkansas, Fayetteville, AR, USA

²Research Division, Arkansas Game and Fish Commission, Little Rock, AR, USA

Correspondence

Tyler K. Chafin, Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO 80309, USA.
Email: tylerkchafin@gmail.com

Present address

Tyler K. Chafin, Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, CO, USA

Funding information

XSEDE, Grant/Award Number: TG-BIO160065; National Science Foundation, Grant/Award Number: DBI-2010774; Arkansas and the Wildlife and Sport Fish Restoration Program, Grant/Award Number: AR-W-F15AF00896; Pittman-Robertson Wildlife Restoration Grant Program, Grant/Award Number: AR-W-F18AF00020

Abstract

Approximately 100 years ago, unregulated harvest nearly eliminated white-tailed deer (*Odocoileus virginianus*) from eastern North America, which subsequently served to catalyze wildlife management as a national priority. An extensive stock-replenishment effort soon followed, with deer broadly translocated among states as a means of re-establishment. However, an unintended consequence was that natural patterns of gene flow became obscured and pretranslocation signatures of population structure were replaced. We applied cutting-edge molecular and biogeographic tools to disentangle genetic signatures of historical management from those reflecting spatially heterogeneous dispersal by evaluating 35,099 single nucleotide polymorphisms (SNPs) derived via reduced-representation genomic sequencing from 1143 deer sampled statewide in Arkansas. We then employed Simpson's diversity index to summarize ancestry assignments and visualize spatial genetic transitions. Using sub-sampled transects across these transitions, we tested clinal patterns across loci against theoretical expectations of their response under scenarios of re-colonization and restricted dispersal. Two salient results emerged: (A) Genetic signatures from historic translocations are demonstrably apparent; and (B) Geographic filters (major rivers; urban centers; highways) now act as inflection points for the distribution of this contemporary ancestry. These results yielded a statewide assessment of contemporary population structure in deer as driven by historic translocations as well as ongoing processes. In addition, the analytical framework employed herein to effectively decipher extant/historic drivers of deer distribution in Arkansas is also applicable for other biodiversity elements with similarly complex demographic histories.

KEYWORDS

ddRADseq, gene flow, genetic drift, historical population dynamics, management, population connectivity, population structure, SNPs, wildlife

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *Evolutionary Applications* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

Understanding movement behaviors and patterns of dispersal across the landscape are important components of a wildlife management strategy. Quantifying dispersal and connectivity among populations has traditionally been done using direct methods such as radio- and satellite-tracking (Kays et al., 2015), yet these methods are labor-intensive, generally only allow for small sample sizes with results difficult to extrapolate at the population level (Katzner & Arlettaz, 2020). An ability to do so indirectly via patterns measured from genetic data such as gene flow (i.e., corresponding to migration, demographics) and genetic drift (i.e., effective population sizes, inbreeding) has made it possible to more accurately assess movement and landscape ecology patterns at a fraction of the per-individual cost (Bossart & Pashley Prowell, 1998; Comte & Olden, 2018; Lowe & Allendorf, 2010). Continual decline of sequencing costs concomitant with computational and analytical advancements now offer an unprecedented density of sampling and geographic resolution that can link individual movement patterns with landscape genetic associations (Richardson et al., 2016), as well as functional genomic correlations (Schoville et al., 2012). This has spurred a proliferation of approaches, such as deducing natal assignment (Battey et al., 2020a), inferring the directionality and pathways of range expansions (Rengefors et al., 2021), and predicting disease spread from spatial host-pathogen associations (Fountain-Jones et al., 2021; Kozakiewicz et al., 2018).

However, contemporary evaluations examine but a single “snapshot” from an evolutionary continuum of events that have cumulatively shaped contemporary diversity, a consequence of which is that genetic patterns may not be interpretable when juxtaposed against modern landscapes (Douglas et al., 1999). Thus, models that invoke landscape genetic associations may incorporate implicit assumptions that lack biological reality (Cushman & Landguth, 2010). One common assumption involves dispersal among a series of discontinuous subpopulations (Lundgren & Ralph, 2019; Petkova et al., 2015) that, when violated, may lead to incorrect inference of discrete causes (such as barrier effects or reproductive isolation; Bradburd et al., 2018). Likewise, gene flow and genetic drift are often conflated, in that one may generate spurious signals of the other (Battey et al., 2020b; Mazet et al., 2016). For example, demographics with a history of turbulence may inflate genetic divergences such that population structure appears artificially discrete and thus suggests restricted gene flow (Austerlitz et al., 1997; Excoffier & Ray, 2008; Nei et al., 2014). A classic example of this phenomenon is the re-expansion of populations following a historical ebb. Upon secondary contact, unique genetic changes that have accumulated during periods of isolation may seemingly initiate an apparent rapid shift in ancestries when viewed from a landscape genetic context (Zellmer & Knowles, 2009).

More recent events such as anthropogenic habitat fragmentation may also act to obscure genetic patterns across the landscape (Epps et al., 2013; Landguth et al., 2010; Pavlacky et al., 2009). In particular, human-mediated dispersal introduces both diversifying

and homogenizing effects that prove difficult to disentangle (Epps & Keyghobadi, 2015; Seebens et al., 2017). This erodes adaptive and/or spatial associations in that it disturbs the expected relationships between genetic differentiation and geographical or environmental proximity (Capinha et al., 2015; Einfeldt et al., 2020). Well-intended actions, such as translocations and stockings, may also inadvertently obscure pre-existing landscape genetic patterns (Brown et al., 2009; Laikre et al., 2010; Shephard et al., 2013). This is especially so for heavily managed species that incur translocations as one aspect of their management (Jahner et al., 2019; Seddon et al., 2014).

White-tailed deer, one of the most recreationally important species in North America, is a case in point (Knoche & Lupi, 2012), with populations intensely impacted by both hunter harvest and game management (Waller & Alverson, 1997; Wolverson et al., 2007). Early in the 20th century, deer and other extensively hunted species (e.g., wild turkey) saw widespread extirpations. In a census of the mid-western United States, Leopold (1931) noted that extant deer populations were of low abundance in southern Missouri, an observation that dovetails with estimated numbers of deer in Arkansas during that period (~500 scattered across isolated refugia at the low ebb; Holder, 1951). Subsequent efforts to bolster resident numbers and re-populate depleted areas in Arkansas involved translocations from within- and out-of-state. Although successful, these efforts promoted genetic patterns within the state that were based on artificial, rather than natural, movements.

Distributional patterns in white-tailed deer clearly reflect multiple factors. Those extrinsic may include rivers, interstate highways, suitable habitat (Robinson et al., 2012); agricultural land use and urbanization (Kelly et al., 2014); and regional eco-physiographic boundaries (Miller et al., 2020). Intrinsic factors, on the other hand, may point to population density (Lutz et al., 2015); age structure/sex ratio (Long et al., 2008); and social hierarchy (Nixon & Mankin, 2016).

A first step in designating factors that have shaped deer population structure and dispersal is to estimate the genetic component within Arkansas deer genomes that still persists as a residual of historic translocations and subsequent genetic drift. Perhaps our most pressing need with regard to dispersal is the impact these data may have in relation to a widespread and fatal neurodegenerative disease of cervids (chronic wasting disease; CWD), which now represents a panzootic (Escobar et al., 2020; Mawdsley, 2020; Williams & Young, 1980). Currently (as of January 2021), the distribution of the disease in Arkansas is limited to a block of counties in the northwest section of the state (Figure 1). Thus, efforts to contain and mitigate its spread are paramount for wildlife management, not only in North America, but also globally (Leiss et al., 2017).

1.1 | Separating contemporary dispersal from historical translocation

The driving factors for deer dispersal primarily relate to habitat quality and climate, and these in turn elicit obvious management interest. Region-specific patterns are needed so as to develop broader




FIGURE 1 Ancestry proportions for $N = 1143$ Arkansas white-tailed deer, as inferred using the program ADMIXTURE applied to 35,099 single-nucleotide polymorphisms (SNPs) generated via ddRAD-seq. Samples are represented as pie charts plotted at absolute collection coordinates, with colors of assignment probabilities proportional to a particular subpopulation. The Arkansas CWD Management Zone (as of January 2021) is indicated in red

generalizations regarding species-specific movement ecology (Brinkman et al., 2005) and to evoke management strategies that can optimize hunter harvest while maintaining population densities.

The goal of our study was to ascertain whether signatures of historic translocations are apparent in, and have contributed to, the genetic diversity and structure of deer in Arkansas, and furthermore, if they can indeed be parsed from a confusing matrix of ongoing gene flow and genetic drift. To do so, we surveyed a broad array of nuclear genomic markers using ddRAD sequencing (Peterson et al., 2012). Our first objective was to characterize respective patterns of anthropogenic translocation and natural re-colonization via refugial populations and secondarily to seek evidence for geographic barriers throughout the state that seemingly prevent or actively filter deer dispersal. However, we hypothesized that the former situation would obscure the latter, in that long-range anthropogenically mediated displacement of individuals violates methodological assumptions that stipulate gene flow as occurring in a spatially consistent manner (Bradburd & Ralph, 2019).

Here, we used ancestry assignment probabilities (e.g., for K number of populations) and reduced this to a single index (Simpson, 1949) that encapsulated the diversity of ancestries within a given spatial grain. This served to reduce those genetic ancestry components that are influenced by widespread translocation, to a single interpolated “surface” representing only changes in ancestry over space (i.e., boundaries between contemporary populations). We also wished to define the respective roles played by stochastic and deterministic processes in generating these zones, particularly given historic population fluctuations and the internal/ external translocations that conflated the demography of Arkansas deer. To do so, we borrowed components of cline theory to hypothesize how individual loci should transition across these spaces relative to a genome-wide average (Barton, 1983; Barton & Hewitt, 1985; Endler, 1973; Hewitt, 2001; Polechová & Barton, 2011; Slatkin, 1973).

Although clinal variation is often taken as evidence for selection (i.e., migration-selection balance; Haldane, 1948), genetic drift in concert with spatially variable gene flow can generate patterns in individual loci that mimic those seen in adaptive clines (Vasemägi, 2006). With drift operating alone, allele frequencies at each locus may wander “upward” or “downward” across space as a function of the initial allele frequency. Range expansion or dissemination from refugia can also yield a type of “rolling” founder effect that can mirror a clinal pattern (Excoffier et al., 2009; Hallatschek & Nelson, 2008; Hewitt, 2000; Keller et al., 2013). Accordingly, we might predict that multi-locus patterns at population boundaries generated solely due to genetic drift in expanding populations will be dominated by stochastic directional change, with each fluctuating either above or below the genome-wide average (Santangelo et al., 2018).

On the other hand, a strong barrier to movement would yield a discontinuity in the rate of directional change across space (Barton, 2008; Nagylaki, 1976; Slatkin, 1973). The existence of a barrier (either impacting dispersal or as a strong fitness differential) thus implies the presence of an inflection point that would induce a rapid shift in locus-specific ancestries (the “width” of this cline is the

inverse of the slope at the inflection point; Endler, 1977; Fitzpatrick, 2013; Slatkin, 1973).

Given these predictions, we summarized locus-wise patterns using two clinal parameters: α , which describes the direction of genetic change, and β , which indicates “width” of a cline. We examined how these two parameters varied, then developed hypotheses regarding the manner by which it impacted our observed population structure. We then considered the evolutionary implications, and limitations, of these results in the context of management, and in so doing demonstrated how the methodologies employed herein can be used to make inferences in other heavily managed species regarding demographic factors (i.e., sex-biased and natal dispersal) and those more extrinsic (i.e., landscape resistance) that drive population structure.

2 | METHODS

2.1 | Sampling and data collection

During 2016–2019, 1720 tissues were collected by the Arkansas Game and Fish Commission (AGFC), representing all 75 Arkansas counties. We employed a combination of targeted sampling, road-kill surveys, and a voluntary statewide CWD testing program (Chafin et al., 2020). Age and sex were collected where possible, with the former estimated by tooth development and wear (Severinghaus, 1949). Data for an additional 30 samples were also attained from Wisconsin to test for signals of historically recorded translocation efforts involving this stock (Holder, 1951). From these, a subset of 1208 samples were chosen for sequencing.

We homogenized tongue or ear tissue (stored at -20°C) and extracted genomic DNA using QIAamp Fast Tissue kits (Qiagen, Inc), with verification via gel electrophoresis (2% agarose). Samples with sufficient yields of high molecular weight DNA (>200 ng) were then enzymatically fragmented via incubation at 37°C , using high-fidelity *NsiI* and *MspI* restriction enzymes (New England Biolabs, Inc.), following enzyme and size-selection optimization using *in silico* digests (Chafin et al., 2018) of several available reference genomes hosted by NCBI: *Odocoileus virginianus* (GCA_002102435.1), *Capreolus capreolus* (GCA_000751575.1) and *Capra hircus* (GCF_001704415.1).

Digests were purified using Ampure XP beads (Beckman-Coulter, Inc.) and standardized to 100 ng per sample. Unique inline barcodes (Peterson et al., 2012) were then ligated using T4 DNA Ligase (following manufacturer protocols; New England Biolabs, Inc.). Samples were then multiplexed ($N = 48$) prior to automated size selection at 300–450 bp using a Pippin Prep (Sage Sciences). Adapter extension was performed over 12 PCR cycles using TruSeq-compatible indexed primers (Illumina, Inc.) and Phusion high-fidelity *taq* polymerase (New England Biolabs, Inc.). Additional quality controls (e.g., qPCR and fragment analysis) were performed on final libraries prior to 1×100 single-end sequencing on the Illumina HiSeq 4000 (Genomics and Cell Characterization Facility, University of Oregon/Eugene), with a total of $N = 96$ samples pooled per lane.

Raw reads were demultiplexed using the pyRAD pipeline (Eaton, 2014), and those with barcode mismatches were discarded. Demultiplexed reads were further filtered by removing those having >4 nucleotides below a quality threshold of 99% accuracy. Reads were then clustered into putative loci within individuals, allowing for a maximum distance threshold of 15%. This was done using the VSEARCH algorithm (Rognes et al., 2016) as implemented in pyRAD, so as to remove read clusters with 3+ indels, >5 ambiguous consensus nucleotides, or a coverage <20X or >500X. Putative homologs were identified using among-individual clustering with the same parameters, and additional removal of loci having >2 alleles per individual, >70% heterozygosity for any polymorphic site, >10 heterozygous sites, or <50% individual recovery (see github.com/tkchafin/scripts for postprocessing and file formatting scripts). To mitigate issues of independence among sites, we additionally sub-sampled the dataset to one SNP per locus.

2.2 | Derivation of population structure

Given known issues with respect to assignment accuracy in datasets dominated by low-frequency variants (Linck & Battey, 2019), we excluded SNPs exhibiting a minor allele count <2 which corresponds to a global minor-allele frequency of ~1%. This minor-allele count threshold per locus (i.e., accounting for total number of sampled alleles), equates to a mean minor-allele frequency threshold of 11% ($SD = 2\%$). We then inferred population structure (ADMIXTURE: Alexander et al., 2009) with parallel processing (ADMIXPIPE: Mussmann et al., 2020). Model selection (i.e., for K , the number of populations) followed a cross-validation approach with results aggregated from 20 independent replicates (CLUMPAK: Kopelman et al., 2015).

Individual-level ADMIXTURE results were summarized as a “surface” with spatial discontinuities represented as interpolated assignment probabilities. Here, we constructed statewide rasters, as representing per-pixel probabilities or “ancestry proportions,” using Empirical Bayesian Kriging (ARCMAP 10.7.1, Esri, Inc.). Probability surfaces were then summarized as evenness and diversity of ancestries in each cell using Simpson's index (Simpson, 1949) (where K =number of statewide subpopulations). Our use of the diversity index was based on a straightforward prediction: Areas representing spatial transitions between populations will have a correspondingly low certainty of assignment to a given subpopulation ancestry (i.e., high diversity). Likewise, those with low inter-population exchange will comprise genetically similar individuals assigned with high probability to the endemic ancestry (i.e., low diversity). Thus, genetic edges represent a marked transition from one population to another, as identified by site-wise diversity in assignment probabilities (i.e. “ancestry diversity”). By lowering the resolution of the genetic diversity surface, we effectively decreased the noise introduced by translocation.

We expect spatial variation in ancestry diversity to be inversely proportional to true rates of gene flow, in that this quantity (as well as the ancestry proportions from which it is computed) is a product

of gene flow averaged over many generations. Of note is the fact that the method only examines local ancestry probabilities and consequently will not be dominated by translocation-related artifacts (as are methods based on interpolated pairwise genetic distances or F_{ST}). Patterns based on Simpson's diversity were contrasted with those inferred using a form of 2D-stepping stone model (EEMS; Petkova et al., 2015), as run with 2 million MCMC iterations (1 million as burn-in), and sampled every 1000 iterations [following parameter sweeps that tuned MCMC acceptance rates to fall between 20% and 50% (Roberts et al., 1997)].

2.3 | Estimating cline parameters

We examined the nature of population boundaries by examining how individual loci transitioned across these regions, relative to the genome-wide average. To do so, we defined eight transects across putative genetic edges, sampling 32–73 individuals per transect ($\bar{x} = 55.38$). Individuals were chosen to represent each subpopulation using a probability threshold applied to ADMIXTURE results. Loci were filtered to remove all SNPs with missing data in >50% of individuals, and computational time was further restricted by retaining only those loci with a sufficient allele frequency differential among clinal extremes [computed as $\delta > 0.50$; (Gregorius & Roberds, 1986)]. Locus-wise clinal patterns were then inferred using a Bayesian method developed originally for hybrid zones (BGC: Gompert & Buerkle, 2011, 2012). Open-source Python code for filtering SNP matrices and generating necessary input is available at (github.com/tkchafin/scripts/phylicp2introgress.pl and [phylicp2bgc.pl](https://github.com/tkchafin/scripts/phylicp2bgc.pl)).

Analyses were performed for each transect across four replicates, each using one million MCMC iterations, discarding the first 500,000 as burn-in, with output thinned to every 500 iterations. Results were summarized by visualizing a 2-D density of cline shape parameters. These are α (= cline directionality) that describes an increase ($\alpha > 0$) or decrease ($\alpha < 0$) in the probability of locus-specific ancestry from a parental population; and β (= cline width/ steepness) that defines the rate of transition in probabilities of locus-specific ancestries having either steep ($\beta > 0$) or wide ($\beta < 0$) shapes (Gompert et al., 2012). In this context, a locus which does not deviate from the genome-wide pattern would have $\alpha \cong \beta \cong 0$. Deviation in a directional manner (i.e., an increase or decrease of one ancestry over another) is described by α , whereas deviation in the rate of ancestry change around an inflection point (i.e., yielding a sigmoidal shape) is described by β . Statistical outliers were designated using the method of Gompert and Buerkle (2011). BGC results were parsed and visualized using the ClineHelpR R package (github.com/btmartin721/ClineHelpR).

2.4 | Relative dispersal by age and sex

Of particular interest in wildlife management is the backward inference of geographic positioning from genotypes—that is, the

Assignment probability

0.00 - 0.20 0.21 - 0.40 0.41 - 0.60 0.61 - 0.80 0.81 - 1.00




FIGURE 2 Assignment probabilities for eight populations of Arkansas white-tailed deer ($k = 1$ through $k = 8$), as inferred from ADMIXTURE applied to 35,099 SNPs, interpolated using Empirical Bayesian Kriging of assignments for $N = 1143$ individuals. $P(k) = 1.0$ corresponds to 100% probability of ancestry per raster cell, and $P(k) = 0.0$ corresponds to 0%. Individual samples are represented as black dots

geolocation of “origination” points for sampled animals. This could be used, for example, to ascertain the geographic origin of poached individuals or to estimate postnatal dispersal. To this end, we used the novel deep-learning method LOCATOR (Battay et al., 2020a) to predict the geographic origin of samples without relying upon explicit assumptions about population genetic processes underlying spatial genetic differentiation (Bradburd & Ralph, 2019). The analysis was performed iteratively across each individual, using the remaining samples to train the LOCATOR classifier, with 100 bootstrap pseudo-replicates to assess variance in geolocation. Given computational constraints, we performed the analysis using a subset of 5000 SNPs having a minor-allele frequency $>10\%$.

We estimated relative dispersal distances as the Euclidean distance between sampled localities and the centroid of predicted coordinates, under the assumption that the distance between predicted and collected locations is the result of lifetime dispersal, at least for samples for which geolocation variance is low among pseudo-replicates. The results were then partitioned by age, sex, and CWD status.

Our second approach examined the decay in genetic relatedness as a function of distance from each individual, measured as a Prevosti distance (R package poppr: Kamvar et al., 2014). Here, the assumption is that recently dispersed individuals will be, on average, more genetically dissimilar from resident individuals, whereas resident individuals having an appreciable reproductive output will be less so. These calculations were limited to individuals that had neighboring samples within a 5 km radius, thus removing individuals from sparsely sampled regions where the relationship between relatedness and distance would be unreliable. We also note that the traditional aging method employed herein (Severinghaus, 1949) has an accuracy seemingly reduced in older deer, potentially suggesting caution in the interpretation of results (Cook & Hart, 1979; Gee et al., 2002; Mitchell & Smith, 1991).

3 | RESULTS

3.1 | Data processing

Our raw data represented $N = 1143$ samples, including $N = 29$ from Wisconsin (Table S1). We removed $N = 83$ that had missing metadata,

discrepancies with coordinates, or $<50\%$ of loci present. Assembly in pyRAD yielded an average of 25,584 loci per sample ($\sigma = 8639$). After removing loci present in $<50\%$ of samples and excluding those containing potential paralogs (e.g., excessive heterozygosity or >2 alleles per locus), our final dataset contained 35,420 loci, from which 2,655,584 SNPs were catalogued. Of these, 54,102 were excluded as singletons. To limit signal redundancy, we then condensed the data to one SNP per locus, yielded a final matrix of 35,099 SNPs for analyses of population structure.

3.2 | Population structure and “ancestry surfaces”

Cross-validation, performed on $N = 20$ replicates each for subpopulation model ($K = 1-20$), revealed the optimal number of clusters as $K = 8$. Spatial orientation of these samples (Figure 1) provided a geographic definition, with some subpopulations qualitatively defined by apparent landscape features, such as the Arkansas River Valley as the southern extent of subpopulations $k3$ and $k6$.

Two ancestries ($k1$ and $k8$) largely dominated south of the Arkansas River, bounded by Interstate 30 to the north and the Ouachita River to the south (Figures 1 and 2), each of which supports the argument that genetic structure is defined by large geographic barriers. The southwestern portion of the state has two ancestral assignments ($k1$ and $k3$), with the latter having mixed representation in the north-central section (potentially an artifact of weak differentiation rather than true shared ancestry). The southeastern section is dominated by a single gene pool ($k8$), which coincidentally subsumed all Wisconsin samples, a strong signature of genetic variability as a residual of historic translocations.

A greater amount of locally endemic structure occurred north of the Arkansas River in the Ozark Mountains, where six subpopulations were evident. The most broadly distributed ($k5$) was to the east in the Mississippi alluvial plains, extending westward across the mainstem of the White River then northward toward the confluence of the Black and White rivers, where it grades into several distinct yet loosely defined subpopulations (Figures 1 and 2). The northwestern corner of the state was the most heterogeneous, with four primarily endemic gene pools ($k = 2, 3, 4$ and 7 ; Figure 2). The

northern-most of these was approximately bounded by the White River (Figure 2) and graded westward into an area with high levels of mixed assignment (Figure 1). The remaining northwestern region was defined by several gene pools showing spatially weak transitions, suggesting reduced gene flow but with geographic and/ or environmental boundaries reasonably porous.

Effective migration surfaces (EEMS) failed to capture any discernable pattern relating to spatially defined population structure (Figure S1). Geographic breaks separating subpopulations (= genetic edges) were captured instead by reducing interpolated assignment probabilities ($n = k$) as a continuous Simpson's diversity index ($n = 1$; Figure 3). This, in turn, reflects a dependence on homogeneity of local assignments, rather than global patterns compounded by long-distance transplants.

3.3 | Intraspecific genomic clines

Genomic clines varied substantially among transects (Figures 4 and S2). Most inter-population comparisons within northwest Arkansas, to include $k2 \times k3$ (both eastern and southern transition zones), $k2 \times k6$, and $k4 \times k7$, indicated variation primarily restricted to cline directionality (α), with cline steepness (β) at a minimum. The variation in locus-wise pattern for these cases (hereafter termed " α -dominant") indicated a directional change in the representation of

reference populations across the transect, but without a noticeable "inflection" point, as implied by nonzero cline width (β).

Remaining comparisons, including one additional transect from northwest Arkansas ($k3 \times k4$), primarily reflected variation in cline width (β), in that loci varied most prominently regarding steepness of transition around an inflection point respective to genome-wide ancestry. Although two transects showed minor exception (i.e., $k3 \times k6$ varied along both parameters; $k4 \times k7$ each showed minimal variation), the contrasting variation between α -dominant versus β -dominant transects suggests that different processes underlie ancestry transitions.

3.4 | Estimating dispersal using geolocation analysis

Individuals from densely sampled regions could be assigned to geographic origin using the "deep-learning" approach, with CWD-positive individuals assigned with a mean bootstrap distance from centroid prediction generally <15 km (Figure S3). However, we did find assignment error (e.g., among bootstraps) was elevated in low-density sampling regions (Figures 5 and S4), which resulted in higher estimated individual dispersal distances (Figure S4A,B). Given that variance in dispersal estimates dropped considerably below ~ 25 km (Figure S4C), a conservative threshold of 10 km was chosen and all individuals having a mean bootstrap-centroid distance above that




FIGURE 3 Simpson's diversity of interpolated ancestry proportions for $N = 1143$ Arkansas white-tailed deer. Each raster cell was assigned eight values equal to the expected proportion of ancestry corresponding to the eight genetic clusters (Figure 1), then summarized using Simpson's diversity index




FIGURE 4 Relationships between genomic cline parameters contrasted among eight sub-sampled transects of individuals spanning population boundaries in Arkansas white-tailed deer (Figure 3). Contour plots show relative densities of SNPs varying in cline steepness (β), representing the gradient of the clines, and cline directionality (α), representing bias in SNP ancestry. Outlier loci are highlighted in blue

were removed for the purposes of dispersal estimates. After filtering, $N = 110$ samples remained (Figure 5A). A higher error threshold (i.e., 20 km) allowed a greater number of total samples to be incorporated ($N = 264$; Figure 5B), with several “roadkills” appearing as long-distance transfers. Although these results are low-precision assignments, they do underscore the capacity of the method regarding the identification of transported individuals (e.g., poached or illegally dumped deer, or carcasses transported across state lines or regional management zones).

Geo-located results for individuals passing a strict error filter demonstrated a dispersal distance for males approximately double that of females across all age classes (statistically significant only for the Y2-2.5 class due to low sample sizes; Figure 6). This pattern was

established as early as the Y1-1.5 group, indicating apparent male dispersal by that age. Smaller dispersal distances were found for fawns across both sexes (Figure 6), again corroborating *a priori* biological expectations.

Patterns of genetic dissimilarity also showed an age x sex effect, with greater genetic dissimilarity for neighboring Y0-1 males than females, and with a shift toward reduced dissimilarity in males >5 (Figure S5). This again supports the argument that male deer in Arkansas have dispersed by the Y1-1.5 age class and potentially reflects age-biased reproduction. Males contributed disproportionately to their local gene pools by age 5 (i.e., producing offspring with resident females), thereby creating a pattern of lower genetic dissimilarity among neighboring individuals, regardless of distance.




FIGURE 5 Summary of geo-location predictions in LOCATOR as a random subset of $N = 490$ Arkansas whitetailed deer at two error thresholds: 10 km (=a) and 20 km (=b), thus constraining results to $N = 110$ (=a) and $N = 264$ (=b). Error calculated as mean distance of bootstrap predictions from the centroid predicted for each individual (see Figure S4). A black dot denotes the predicted location of an individual and a colored dot indicates the ‘true’ location (colors proportional to measurement error in km)




FIGURE 6 Inferred dispersal distances, partitioned by sex and age class for $N = 1143$ Arkansas white-tailed deer. Dispersal distances were calculated as the difference (in km) between the predicted and true locations, excluding all individuals with a mean prediction error of 10 km. Sample sizes for each group are given in black, with the mean in red below each box plot. Within-age, p-values are reported for two-sample t-tests comparing means from males and females




FIGURE 7 Proposed translocation pathways and historical refuge populations for Arkansas white-tailed deer, showing (left) translocation events supported by genetics along (dashed line), as well as those with anecdotal support given limited historical records (solid), from three primary stocking sources: Sandhill Game Farm in Wisconsin (W); ‘Sylamore District’ state refuge sites (S); and the Howard County Game Farm (H). Also shown are major inhabited regions of white-tailed deer (right) estimated from 1942–47 game surveys, excluding small regions (see full version in Online Supplementary Material. Putative inhabited regions are also annotated according to their hypothesized associations with contemporary genetic clusters (Figures 1 and 2)

4 | DISCUSSION

4.1 | Genetic footprints of historic management

The prolonged history of hunter harvest, and subsequent long-range translocations into Arkansas and the surrounding region (Ellsworth et al., 1994; Holder, 1951), provide a sideboard to our study in that artificial long-distance movements such as these violate the assumptions inherent with many spatially explicit methods. However, we

found estimates of ancestry diversity and probability surfaces to be qualitatively robust in that they firmly recapitulated the record of historic translocations (Holder, 1951). This, in turn, necessitates that our results be placed within an historic context.

In the early 20th century, following many decades of over-hunting, the Arkansas deer population declined to <500 individuals (Holder, 1951). In response, Arkansas Game and Fish Commission (AGFC) implemented an extensive restocking program (1941–1951) involving as its basis three primary sources. The first was Howard

County deer farm (southwestern Arkansas; Figure 7), established from locally transplanted central Arkansas individuals (Wynn, 1943), and now located within the epicenter of population k1 (Figures 2 and 7). Its ancestry is shared elsewhere in the state, thus establishing it as an epicenter for local translocations.

A second major source was the Sylamore District in the Ozark Mountains (Holder, 1951; Wood, 1944), where individuals were naturally abundant (Figure S7). Individuals from this region have a consistently higher probability of assignment to k3 (Figures 2 and 7), as records indicate ~81% of repopulated individuals in the Gulf Coastal Plain originated from there (Holder, 1951; Karlin et al., 1989; Wood, 1944). Our analyses agree that many individuals from southwestern Arkansas (Figures 1 and 6) reflect a mixed assignment to the "Sylamore" cluster. They also comprised ~36% of stocking efforts in the Mississippi Delta region (Karlin et al., 1989).

Our results also indicate some mixed assignment of individuals to k3, although a more widespread representation is found in k5. AGFC surveys (~1942–1945) indicated deer as being relatively abundant in the southeastern Delta region (Figures 7 and S7), seemingly linked with re-population efforts following the disastrous 1927 Mississippi flood. This event, coupled with over-hunting, nearly extirpated deer in the region, save small numbers sustained by local sportsmen (Holder, 1951; Figure S7). We hypothesize deer in the Delta region largely result from those efforts, given the contemporary homogeneity of ancestry assignments in this region.

A third (extraneous) source was the Sandhill Game Farm (Babcock, WI; Wood, 1944). Records indicate that ~64% of deer released into the Mississippi Delta region originated out-of-state, the majority from Wisconsin (Holder, 1951; Karlin et al., 1989). Our Wisconsin samples were unanimously assigned to a gene pool prominently represented in the southern Delta region (k8; Figure 2), firmly establishing its genetic legacy as extending from imported deer. Mitochondrial haplotypes putatively originating from Wisconsin were also uncovered in Missouri, Kentucky, and Mississippi (Budd et al., 2018; DeYoung et al., 2003; Doerner et al., 2005).

All genetic "clusters" that lacked spatial cohesion in our analyses can be connected to the three major stocking sources involved in earlier restoration efforts (per historic records). The remaining sub-populations, primarily in the Ozarks (k2, 4, 6, and 7), may represent natural re-colonization from refugial populations (Figures 2 and 7), a hypothesis supported by early census data (Figures S7 and S8). If this was not recognized, then any analysis of deer genetic structure across Arkansas would instead infer unusually high rates of gene flow among these geographically distant regions. This, in turn, would preclude any analysis aimed at linking variation in genetic structure to environmental factors.

4.2 | Contemporary genetic structure

A primary focus in our study was the contemporary genetic structure of Arkansas deer, especially regarding how these data promote an understanding of landscape resistance to dispersal

(Hemming-Schroeder et al., 2018; Kelly et al., 2014). Yet several prerequisites are apparent in this regard. For example, one apparent question previously constrained by technological limitations is the degree to which potential patterns of genetic variability in deer have been conflated by anthropogenically mediated translocations.

We addressed this issue by utilizing next-generation molecular techniques to derive highly variable markers across the genome. We also implemented advanced geospatial procedures that visualized the spatial transitions inherent within deer ancestry in such a way that translocation histories were not a limiting factor. We did so by interpolating our assignment probabilities from ADMIXTURE, then applying Simpson's diversity index as a means of reducing the probabilities to a single vector. We also polarized stochastic *versus* deterministic processes by examining locus-specific patterns within our transitions.

Previous studies concluded that inferences from landscape genetic analyses were limited at best, due to complex interactions between historic translocations and subsequent population growth (Budd et al., 2018; Leberg et al., 1994; Leberg & Ellsworth, 1999). However, many of these studies relied upon legacy molecular markers (e.g., mtDNA or reduced panels of microsatellite DNA markers) that capture substantially less polymorphism than do next-generation methods (Hodel et al., 2017; Jeffries et al., 2016; Lemopoulos et al., 2019). Recent studies at more refined spatial scales have supported large-scale geographic barriers (e.g., rivers, highways) as being semi-permeable to gene flow (Kelly et al., 2014; Locher et al., 2015; Miller et al., 2020; Robinson et al., 2012). This result is in accordance with radio-telemetry data (Peterson et al., 2017).

Several scale-dependent issues may modulate the degree to which translocations obfuscate landscape genetic patterns. First, as spatial scale increases, so does the probability of sampling individuals whose genetic dissimilarity reflects translocation rather than natural gene flow via dispersal. Second, the most obvious environmental effects that drive deer dispersal have the greatest probability of being recovered in an analysis, even if they represent spurious signals driven by translocation. However, landscape-level analyses of dispersal can still be informative, despite those rapid transitions between historic and contemporary conditions (Epps & Keyghobadi, 2015) that may obfuscate relationships. For example, large-scale environmental features also drive spatial genetic patterns in invasive species, despite reduced levels of genetic diversity (Lopez et al., 2014; Sacks et al., 2016; Zalewski et al., 2009). Such patterns are also rapidly manifested within populations occupying habitats recently modified by anthropogenic developments, such as large urban centers (Beninde et al., 2016; Combs et al., 2018; Kimmig et al., 2020).

The analytical artifacts derived from translocation are expected to be not only scale-dependent but also a function of specific analytical assumptions. For example, classic tests of "isolation-by-distance" (IBD) implicitly assume a negative relationship between geographic proximity and pairwise patterns of genetic distance (Meirmans, 2012; Rousset, 1997). Yet, translocations are demonstrably inconsistent with this expectation. Introduced individuals reflect levels of genetic similarity that are demonstrably inconsistent with spatial

and/ or environmental distance (i.e., “resistance”) that now separates them. This, in turn, incorporates spurious signals within the landscape inferences subsequently derived.

The assignment method used in our study yielded patterns of genetic similarity that reflected translocations as depicted in historic records (Figure 7). We sought to more appropriately expose landscape features that could potentially modulate deer dispersal by effectively “removing” this artificial signal. Our approach (i.e., reducing the dimensionality of assignment probabilities) produced relatively straightforward predictions about the ensuing metric: Areas with uninterrupted movement of individuals should have increased homogeneity regarding interpolated ancestry assignments. By the same logic, regions representing barriers to movement among regions should likewise demarcate transitions in interpolated ancestries. We acknowledge that spatial assignments can also be vulnerable to artifacts, such as an over-fitting of discrete clusters within otherwise continuous populations (Bradburd et al., 2018). Yet despite this, they are demonstrably effective at identifying linear barriers to gene flow (Blair et al., 2012). Coincidentally, our approach using summarized ancestries revealed numerous linear subdivisions that aligned with major landscape barriers, such as rivers in Arkansas (Figure 3). Other recognized transitions approximately corresponded with large urban centers. Our argument is that approaches involving an assumption of stepwise gene flow (implicit with pairwise genetic distances) may generate conclusions that either lack the necessary nuances or are altogether incorrect. This is apparent when the “surfaces” inferred using Simpson’s diversity of ancestry assignments (Figure 3) are contrasted with EEMS results (Figure S1).

4.3 | Separating stochastic versus deterministic processes in clinal patterns

When deriving clinal patterns, it becomes difficult to disentangle contemporary demographic processes from analytical idiosyncrasies. As a means of clarification, we generated multi-locus genomic patterns under the expectation that the relative importance of stochastic versus deterministic processes could be derived from the manner by which genome-wide averages deviated from locus-wise clines (Barton, 1983; Barton & Hewitt, 1985; Slatkin, 1973; Vasemägi, 2006). We found eight transects that varied either in the steepness of genetic transitions (β), or the directionality of allelic frequency change (α). Interestingly, “ α -dominant” transects were found within the Ozark region, previously hypothesized as being naturally re-colonized from local refugia (i.e., $k2 \times k3E/S$, $k2 \times k6$, and $k4 \times k7$; Figures 4 and S2).

Under this hypothesis, genetic drift along the leading edge of population expansions would generate two genetic reverberations: The random, directional fixation of alleles, and an apparent inflation of population structure. Both were observed in our data (Figures 1 and 4). The hypothesis is also supported by statewide surveys (1940’s) that found deer in habitat associated with each Ozark sub-population during, and immediately following, the low ebb of deer in

Arkansas (Figures 7 and S8). The transects likewise displayed near-zero β values, suggesting the absence of a “true” barrier to dispersal. Here, the expectation would be that a hard barrier between discrete populations would reflect changes in allele frequencies around an inflection point.

One interpretation is that genetic drift at the edge of an expanding refugium creates a “rolling” founder effect, with alleles either over- ($\alpha > 0$) or under-represented ($\alpha < 0$) across the re-colonized territory (Excoffier & Ray, 2008; Hallatschek & Nelson, 2008). This phenomenon—termed “gene surfing”—can generate wholly neutral patterns that appear to be adaptive (Peischl et al., 2016; Travis et al., 2007) and may explain the presence of numerous α -outlier loci in those transects (Figure 4).

Interestingly, all our β -dominant transects crossed major rivers: Transect $k3 \times k6$ spanned the Arkansas River; $k3 \times k5$ the White River; and $k1 \times k8$ the Ouachita River. Transect $k3 \times k4$ crosses the smaller and shallower Buffalo River, which may explain the depressed variability in cline steepness (Figure 4). The coincidence of these patterns with major rivers suggests barriers to individual movement, in that nonzero β values implicate variability in the “steepness” of clinal transitions around an inflection point [i.e., with rates of change in allele frequencies either steep ($\beta > 0$) or wide ($\beta < 0$)].

Classically, clinal patterns are established either by selection or hard boundaries to gene flow (Endler, 1973; Slatkin, 1973), but our analyses (Figure 4) revealed little evidence of selection-driven outliers. However, spatially variable gene flow, when coupled with drift, can yield clinal patterns across a considerable proportion of the genome without necessitating selection (Vasemägi, 2006). Selection could feasibly be involved, at least for regions where different deer subspecies were translocated, in that subspecific crosses have elicited fitness impacts such as dystocia (abnormal maternal labor due to shape, size, or position of the fetus; Galindo-Leal & Weber, 1994). The genetic costs of inter-subspecific stocking are recognized (Hopken et al., 2015) and manifested as an anomalous variation in breeding time among other southern-recovered deer populations (Summers et al., 2015). Some limited evidence also for reproductive isolation among mule deer subspecies (*Odocoileus hemionus*) as driven by pheromonal differences (Müller-Schwarze & Müller-Schwarze, 1975).

4.4 | Management implications

Our results identified a diagnosable genetic signature of historic translocations within the genomes of extant Arkansas deer populations. This simultaneously underscores the success of early restocking efforts, while also reiterating long-standing concerns about the genetic and/ or phenotypic impacts of anthropogenic translocations (Meffe & Vrijenhoek, 1988). It also reinforces the need to formalize as an explicit management consideration the intraspecific taxonomy of deer, both to better understand the (yet to be seen) evolutionary consequences of past translocations and to provide a more proactive baseline going forward (Cronin, 2003; Gippoliti et al., 2018).

Resource agencies, in particular, should prioritize those populations that retain endemic genetic diversity and, by so doing, acknowledge the importance of preserving evolutionary legacy (Crandall et al., 2000). Such recognition would capitalize on the many years of natural selection that have operated on those populations prior to anthropogenic interference, which may have generated adaptive genetic variation necessary to withstand changing conditions, or mediated an evolutionary response to disease (e.g., in the *PRNP* gene; Chafin et al., 2020). This further emphasizes the importance of integrating genomic methods for heavily managed species (Flanagan et al., 2018), given their recognized lack of intraspecific adaptive variation [i.e., a “Darwinian shortfall” in biodiversity conservation (Diniz-Filho et al., 2013)].

It is noteworthy that the current boundaries for the AGFC “Deer Management Units” (DMUs) (Meeker et al., 2019) are remarkably consistent with many of the population boundaries identified in our study. This validates the continued use of these ecosystem-based management units for the application of locally appropriate management and harvest regimes.

Another valuable application of our results would be disease containment, particularly given that CWD is a significant challenge globally for the management of cervids (Rivera et al., 2019; Uehlinger et al., 2016). The spread of CWD will be more rapid within rather than among populations, and the cumulative risk of disease spread to new populations will depend on both dispersal and disease prevalence. Thus, potential mitigation efforts should work to constrain disease transmission within CWD-affected areas as well as limiting the outward dispersal of individuals from those areas. One approach to reduce dispersal is to reduce the density of yearling males, which often comprise >50% of emigrating individuals (McCoy et al., 2005; Nelson, 1993; Nixon et al., 1991). Our patterns of age- and sex-biased dispersal distances (Figures 5 and 6) largely recapitulate these expectations at a local level. Removal of young males to slow dispersal also may be compatible with the application of male-focused harvest strategies predicted to reduce disease prevalence (Potapov et al., 2016). A similar action would be to reduce local population densities that serve to drive dispersal, in that robust population numbers elevate social pressure as well as resource competition (Long et al., 2008; Lutz et al., 2015; Shaw et al., 2006). Other harvest strategies have been implemented that effectively decrease emigration while focusing on meeting specific demography goals including actively decreasing overall population density, elevating the proportion of mature males in the population, and increasing the female component of the sex ratio (Brothers & Ray, 1975; Hamilton et al., 1995; Long et al., 2008; Lutz et al., 2015; McCoy et al., 2005; Shaw et al., 2006). However, mature male deer tend to demonstrate higher CWD prevalence than other demographic groups, and management strategies that maximize this component of the population may exacerbate disease dynamics (Miller et al., 2020). Additionally, our findings indicate that disease surveillance strategies should include efforts that focus on detecting “breaches” between genetically distinct regions (those bordered by major rivers and urban centers, as herein; Figure 3) to further direct management and response efforts.

4.5 | Utility of genomic approaches for white-tailed deer management

Beyond providing a nuanced view of historical management processes, the genomic methods used herein offer several improvements beyond “legacy” approaches. For example, they provide vastly increased statistical power for individual-level genetic analyses, such as estimating relatedness (Lemopoulos et al., 2019), or parentage assignment (Flanagan & Jones, 2019). This is also accomplished at a fraction of the cost-per-marker (Puckett, 2017). Evidence for this is found in our geolocation results (Figures 5, S3 and S4), where a lower dispersion of predicted natal locations was achieved when compared with previous attempts using microsatellites (Puckett & Eggert, 2016). The genome is also more densely sampled and these data can be used in conjunction with the analytical methods used herein to infer both environmental associations and adaptive genetic variation (Martin et al., 2020), or detect anthropogenically mediated change through time (Chafin et al., 2019).

SNP-based methods have a particular advantage with regard to scalability (i.e., such as in a multi-state consortium of wildlife management agencies). Large datasets (as herein) may allow a subset of highly consistent, maximally informative markers to be extracted that are broadly applicable across states and regions. An added benefit of our ddRAD dataset is that it provides a broad geographic database from future studies that can guide assay design (e.g., Chafin et al., 2018). Because sequencing effort in the method we used (to include other RADseq methods) may be wasted on loci which have either nonrandom distributions of missing data or low information content (Eaton, 2014; Huang & Knowles, 2016; Rubin et al., 2012), future studies can achieve a vastly improved per-sample cost efficiency by targeting a reduced set of maximally informative loci by mining our dataset for use with targeted-enrichment approaches [e.g., Rapture (Hoffberg et al., 2016)] or those amplicon-based [GT-Seq (Campbell et al., 2015)].

For white-tailed deer, our approach provides a broad geographic reference database, followed by assay design for a *post hoc* method (such as that of Campbell et al., 2015). This would allow our methodology to be deployed so as to generate a sampling sufficiently dense for management-oriented population genetic analyses such as natal geolocation. This, in turn, would facilitate the detection of illegal carcasses, deer trafficking, migrant detection, or characterization of historic translocation (as herein). Moreover, such an approach would be fully replicable across laboratories and agencies, thereby facilitating inter-agency and multi-state collaboration.

Although dense spatial and genomic sampling may offer a composite resolution (e.g., Chafin et al., 2020; Linck et al., 2019), it does not necessarily supplant other approaches. For example, sampling density was shown herein to be a particular limiting factor in geolocation analyses, with high prediction error in regions of the state sparsely sampled (Figure 5). Although we did not directly compare our methodology with that based on a microsatellite dataset, we suspect greater error in the latter given decreased information content associated with the much-reduced dataset. Genetic input from

unsampled populations (e.g., long-distance migrants from out-of-state) found at edges of our sampling areas could also inflate assignment uncertainty (see Figure 3), as could extremely uneven sampling (Lawson et al., 2018). We are also unsure of the degree to which long-distance migrants could influence our clinal analyses (e.g., via a “smoothing” effect), nor fluctuation that may occur over brief time scales (as observed in other study systems; Chafin et al., 2019). Thus, we caution that the deployment of these methods at multi-state or regional scales should focus on dense, spatially expansive sampling that employs both spatial and temporal replicates, where possible (Short Bull et al., 2011). Both priorities are facilitated via the aforementioned enrichment methods that drastically reduce per-sample cost efficiency.

ACKNOWLEDGMENTS

The authors thank hunters and wildlife managers statewide who collected WTD tissue samples. We give particular thanks to A.J. Riggs for helping to coordinate sample collection. We also thank L. Graham for contributions to study design, logistics, and DNA extraction, S.M. Musmann for assistance with geospatial analyses, and J.F. Pummil for facilitating access to computational resources. These were provided by the Arkansas High Performance Computing Center via multiple National Science Foundation grants, the Arkansas Economic Development Commission, and an XSEDE Research Allocation (TG-BIO160065) that provided access to the JetStream cloud service. We are also indebted to the University of Arkansas Library Special Collections for access to archived materials relating to the history of wild game in Arkansas. This research project was funded by the State of Arkansas and the Wildlife and Sport Fish Restoration Program (#AR-W-F15AF00896), and also in part by the Pittman-Robertson Wildlife Restoration Grant Program (#AR-W-F18AF00020) of the U.S. Fish and Wildlife Service through an agreement with the Arkansas Game and Fish Commission. Additional funding was also provided by several generous endowments from the University of Arkansas: Distinguished Doctoral Fellowships (TKC, ZDZ), the Bruker Professorship in Life Sciences (MRD), and the Twenty-First Century Chair in Global Change Biology (MED). TKC is additionally funded by an NSF Postdoctoral Fellowship in Biology (DBI: 2010774). The opinions expressed herein do not necessarily reflect the views or policies of the Arkansas Game and Fish Commission, U.S. Fish and Wildlife Service, nor any additional funding sources or affiliated organization. Product references do not imply endorsements.

CONFLICT OF INTEREST

None declared.

AUTHOR CONTRIBUTIONS

All authors contributed to study design and conceptualization. MCG, CRM, and JRB coordinated sample collection and curation of samples; TKC, BTM, ZDZ, and MRD contributed to molecular work; TKC, ZDZ, MRD, and MED planned the analysis approach; TKC and BTM wrote code for analysis and contributed exploratory analyses;

TKC performed bioinformatic work, assembled data, and performed analyses; TKC and ZDZ prepared figures; TKC and ZDZ drafted the manuscript, and all authors revised the manuscript and contributed to the final version.

DATA AVAILABILITY STATEMENT

Raw sequences are accessioned in the NCBI Sequence Read Archive (SRA) under BioProject PRJNA690954; Assembled SNP data and sample metadata are available via the Open Science Framework (<https://doi.org/10.17605/OSF.IO/T82RV>); codes and custom scripts developed in support of this work are also available as open-source via GitHub under the GNU Public License: github.com/tkchafin (and as cited in-text).

ORCID

Tyler K. Chafin  <https://orcid.org/0000-0001-8687-5905>

REFERENCES

- Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19(9), 1655–1664.
- Austerlitz, F., Jung-Muller, B., Godelle, B., & Gouyon, P.-H. (1997). Evolution of coalescence times, genetic diversity and structure during colonization. *Theoretical Population Biology*, 51, 148–164.
- Barton, N. H. (1983). Multilocus clines. *Evolution*, 37(3), 454–471.
- Barton, N. H. (2008). The effect of a barrier to gene flow on patterns of geographic variation. *Genetics Research*, 90(1), 139–149.
- Barton, N. H., & Hewitt, G. M. (1985). Analysis of hybrid zones. *Annual Review of Ecology and Systematics*, 16(1), 113–148.
- Batthey, C. J., Ralph, P. L., & Kern, A. D. (2020a). Predicting geographic location from genetic variation with deep neural networks. *eLife*, 9, e54507.
- Batthey, C. J., Ralph, P. L., & Kern, A. D. (2020b). Space is the place: Effects of continuous spatial structure on analysis of population genetic data. *Genetics*, 215(1), 193–214.
- Beninde, J., Feldmeier, S., Werner, M., Peroverde, D., Schulte, U., Hochkirch, A., & Veith, M. (2016). Cityscape genetics: structural vs. functional connectivity of an urban lizard population. *Molecular Ecology*, 25(20), 4984–5000.
- Blair, C., Weigel, D. E., Balazik, M., Keeley, A. T. H., Walker, F. M., Landguth, E., Cushman, S., Murphy, M., Waits, L., & Balkenhol, N. (2012). A simulation-based evaluation of methods for inferring linear barriers to gene flow. *Molecular Ecology Resources*, 12(5), 822–833.
- Bossart, J. L., & Pashley Prowell, D. (1998). Genetic estimates of population structure and gene flow: Limitations, lessons and new directions. *Trends in Ecology and Evolution*, 13(5), 202–206.
- Bradbud, G. S., Coop, G. M., & Ralph, P. L. (2018). Inferring continuous and discrete population genetic structure across space. *Genetics*, 210, 33–52.
- Bradbud, G. S., & Ralph, P. L. (2019). Spatial population genetics: It's about time. *Annual Review of Ecology, Evolution, and Systematics*, 50(1), 427–449.
- Brinkman, T. J., Deperno, C. S., Jenks, J. A., Haroldson, B. S., & Osborn, R. G. (2005). Movement of female white-tailed deer: Effects of climate and intensive row-crop agriculture. *Journal of Wildlife Management*, 69(3), 1099–1111.
- Brothers, A., & Ray, M. E. (1975). *Producing quality whitetails*. Wildlife Services.
- Brown, S. K., Hull, J. M., Updike, D. R., Fain, S. R., & Ernest, H. B. (2009). Black bear population genetics in California: Signatures of population structure, competitive release, and historical translocation. *Journal of Mammalogy*, 90(5), 1066–1074.

- Budd, K., Berkman, L. K., Anderson, M., Koppelman, J., & Eggert, L. S. (2018). Genetic structure and recovery of white-tailed deer in Missouri. *Journal of Wildlife Management*, 82(8), 1598–1607.
- Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources*, 15(4), 855–867.
- Capinha, C., Essl, F., Seebens, H., Moser, D., & Pereira, H. M. (2015). The dispersal of alien species redefines biogeography in the Anthropocene. *Science*, 348(6240), 1248–1251.
- Chafin, T. K., Douglas, M. R., Martin, B. T., & Douglas, M. E. (2019). Hybridization drives genetic erosion in sympatric desert fishes of western North America. *Heredity*, 123(6), 759–773.
- Chafin, T. K., Douglas, M. R., Martin, B. T., Zbinden, Z. D., Middaugh, C. M., Ballard, J., Gray, M. C., White, D. Jr., & Douglas, M. E. (2020). Age structuring and spatial heterogeneity in prion protein gene (PRNP) polymorphism in Arkansas white-tailed deer. *Prion*, 14(1), 238–248.
- Chafin, T. K., Martin, B. T., Musmann, S. M., Douglas, M. R., & Douglas, M. E. (2018). FRAGMATIC: In silico locus prediction and its utility in optimizing ddRADseq projects. *Conservation Genetics Resources*, 10(3), 325–328.
- Combs, M., Puckett, E. E., Richardson, J., Mims, D., & Munshi-South, J. (2018). Spatial population genomics of the brown rat (*Rattus norvegicus*) in New York City. *Molecular Ecology*, 27(1), 83–98.
- Comte, L., & Olden, J. D. (2018). Fish dispersal in flowing waters: A synthesis of movement- and genetic-based studies. *Fish and Fisheries*, 19(6), 1063–1077.
- Cook, R. L., & Hart, R. V. (1979). Ages assigned known-age Texas white-tailed deer: Tooth wear versus cementum analysis. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies*.
- Crandall, K. A., Bininda-Emonds, O. R. R., Mace, G. M., & Wayne, R. K. (2000). Considering evolutionary processes in conservation biology. *Trends in Ecology and Evolution*, 15(7), 290–295.
- Cronin, M. A. (2003). Research on deer taxonomy and its relevance to management. *Ecoscience*, 10(4), 432–442.
- Cushman, S. A., & Landguth, E. L. (2010). Spurious correlations and inference in landscape genetics. *Molecular Ecology*, 19(17), 3592–3602.
- DeYoung, R. W., Demarais, S., Honeycutt, R. L., Rooney, A. P., Gonzales, R. A., & Gee, K. L. (2003). Genetic consequences of white-tailed deer (*Odocoileus virginianus*) restoration in Mississippi. *Molecular Ecology*, 12(12), 3237–3252.
- Diniz-Filho, J. A. F., Loyola, R. D., Raia, P., Mooers, A. O., & Bini, L. M. (2013). Darwinian shortfalls in biodiversity conservation. *Trends in Ecology and Evolution*, 28(12), 689–695.
- Doerner, K. C., Braden, W., Cork, J., Cunningham, T., Rice, A., Furman, B. J., & McElroy, D. (2005). Population genetics of resurgence: White-tailed deer in Kentucky. *Journal of Wildlife Management*, 69(1), 345–355.
- Douglas, M. E., Minckley, W. L., & DeMarais, B. D. (1999). Did vicariance mold phenotypes of western North American fishes? Evidence from Gila River cyprinids. *Evolution*, 53(1), 238–246.
- Eaton, D. A. R. (2014). PyRAD: Assembly of de novo RADseq loci for phylogenetic analyses. *Bioinformatics*, 30(13), 1844–1849.
- Einfeldt, A. L., Jesson, L. K., & Addison, J. A. (2020). Historical human activities reshape evolutionary trajectories across both native and introduced ranges. *Ecology and Evolution*, 10(13), 6579–6592.
- Ellsworth, D. L., Honeycutt, R. L., Silvy, N. J., Smith, M. H., Bickham, J. W., & Klimstra, W. D. (1994). White-tailed deer restoration to the southeastern United States: Evaluating genetic variation. *The Journal of Wildlife Management*, 58(4), 686.
- Endler, J. A. (1973). Gene flow and population differentiation. *Science*, 179(4070), 243–250.
- Endler, J. A. (1977). *Geographic variation, speciation, and clines*. Princeton University Press.
- Epps, C. W., & Keyghobadi, N. (2015). Landscape genetics in a changing world: Disentangling historical and contemporary influences and inferring change. *Molecular Ecology*, 24(24), 6021–6040.
- Epps, C. W., Wasser, S. K., Keim, J. L., Mutayoba, B. M., & Brashares, J. S. (2013). Quantifying past and present connectivity illuminates a rapidly changing landscape for the African elephant. *Molecular Ecology*, 22(6), 1574–1588.
- Escobar, L. E., Pritzkow, S., Winter, S. N., Grear, D. A., Kirchgessner, M. S., Dominguez-Villegas, E., Machado, G., Townsend Peterson, A., & Soto, C. (2020). The ecology of chronic wasting disease in wildlife. *Biological Reviews*, 95(2), 393–408.
- Excoffier, L., Foll, M., & Petit, R. J. (2009). Genetic consequences of range expansions. *Annual Review of Ecology, Evolution, and Systematics*, 40(1), 481–501.
- Excoffier, L., & Ray, N. (2008). Surfing during population expansions promotes genetic revolutions and structuration. *Trends in Ecology and Evolution*, 23(7), 347–351.
- Fitzpatrick, B. M. (2013). Alternative forms for genomic clines. *Ecology and Evolution*, 3(7), 1951–1966.
- Flanagan, S. P., Forester, B. R., Latch, E. K., Aitken, S. N., & Hoban, S. (2018). Guidelines for planning genomic assessment and monitoring of locally adaptive variation to inform species conservation. *Evolutionary Applications*, 11(7), 1035–1052.
- Flanagan, S. P., & Jones, A. G. (2019). The future of parentage analysis: From microsatellites to SNPs and beyond. *Molecular Ecology*, 28(3), 544–567.
- Fountain-Jones, N. M., Kraberger, S., Gagne, R., Trumbo, D. R., Salerno, P., Chris Funk, W., Crooks, K., Biek, R., Alldredge, M., Logan, K., Baele, G., Dellicour, S., Ernest, H. B., VandeWoude, S., Carver, S., & Craft, M. E. (2021). Host relatedness and landscape connectivity shape pathogen spread in a large secretive carnivore. *Communications Biology*, 4, 12.
- Galindo-Leal, C., & Weber, M. (1994). Translocation of deer subspecies: Reproductive implications. *Wildlife Society Bulletin*, 22(1), 117–120.
- Gee, K. L., Holman, J. H., Causey, M. K., Rossi, A. N., & Armstrong, J. B. (2002). Aging white-tailed deer by tooth replacement and wear: A critical evaluation of a time-honored technique. *Wildlife Society Bulletin*, 30(2), 387–393.
- Gippoliti, S., Cotterill, F. P. D., Groves, C. P., & Zinner, D. (2018). Poor taxonomy and genetic rescue are possible co-agents of silent extinction and biogeographic homogenization among ungulate mammals. *Biogeographia*, 33(1), 41–54.
- Gompert, Z., & Buerkle, A. C. (2011). Bayesian estimation of genomic clines. *Molecular Ecology*, 20, 2111–2127.
- Gompert, Z., & Buerkle, C. A. (2012). bgc: Software for Bayesian estimation of genomic clines. *Molecular Ecology Resources*, 12(6), 1168–1176.
- Gompert, Z., Parchman, T. L., & Buerkle, C. A. (2012). Genomics of isolation in hybrids. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1587), 439–450.
- Gregorius, H. R., & Roberds, J. H. (1986). Measurement of genetical differentiation among subpopulations. *Theoretical and Applied Genetics*, 71, 826–834.
- Haldane, J. B. S. (1948). The theory of a cline. *Journal of Genetics*, 48(3), 277–284.
- Hallatschek, O., & Nelson, D. R. (2008). Gene surfing in expanding populations. *Theoretical Population Biology*, 73(1), 158–170.
- Hamilton, R. J., Know, W. M., & Guynn, D. C. (1995). How quality deer management works. In K. V. Miller, & R. L. Marchinton (Eds.), *Quality whitetails: The why and how of Quality Deer Management* (pp. 7–18). Stackpole.
- Hemming-Schroeder, E., Lo, E., Salazar, C., Puente, S., & Yan, G. (2018). Landscape genetics: A toolbox for studying vector-borne diseases. *Frontiers in Ecology and Evolution*, 6, 1–11.

- Hewitt, G. M. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, 405(6789), 907–913.
- Hewitt, G. M. (2001). Speciation, hybrid zones and phylogeography—seeing genes in space and time. *Molecular Ecology*, 10(3), 537–549.
- Hodel, R. G. J., Chen, S., Payton, A. C., McDaniel, S. F., Soltis, P., & Soltis, D. E. (2017). Adding loci improves phylogeographic resolution in red mangroves despite increased missing data: Comparing microsatellites and RAD-Seq and investigating loci filtering. *Scientific Reports*, 7(1), 1–14.
- Hoffberg, S. L., Kieran, T. J., Catchen, J. M., Devault, A., Faircloth, B. C., Mauricio, R., & Glenn, T. C. (2016). RADcap: Sequence capture of dual-digest RADseq libraries with identifiable duplicates and reduced missing data. *Molecular Ecology Resources*, 16(5), 1264–1278.
- Holder, T. H. (1951). A survey of Arkansas game. *Arkansas Game and Fish Commission Federal Aid Publication Project II-R.*, 57–79.
- Hopken, M. W., Lum, T. M., Meyers, P. M., & Piaggio, A. J. (2015). Molecular assessment of translocation and management of an endangered subspecies of white-tailed deer (*Odocoileus virginianus*). *Conservation Genetics*, 16(3), 635–647.
- Huang, H., & Knowles, L. L. (2016). Unforeseen consequences of excluding missing data from next-generation sequences: Simulation study of RAD sequences. *Systematic Biology*, 65(3), 357–365.
- Jahner, J., Matocq, M. D., Malaney, J. L., Cox, M., Wolff, P., Gritts, M. A., & Parchman, T. L. (2019). The genetic legacy of 50 years of desert bighorn sheep translocations. *Evolutionary Applications*, 12(2), 198–213.
- Jeffries, D. L., Copp, G. H., Lawson Handley, L., Olsén, K. H., Sayer, C. D., & Hänfling, B. (2016). Comparing RADseq and microsatellites to infer complex phylogeographic patterns, an empirical perspective in the Crucian carp, *Carassius carassius*, L. *Molecular Ecology*, 25(13), 2997–3018.
- Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, 2, e281.
- Karlin, A. A., Heidt, G. A., & Sugg, D. W. (1989). Genetic variation and heterozygosity in white-tailed deer in southern Arkansas. *The American Midland Naturalist*, 121(2), 273–284.
- Katzner, T. E., & Arlettaz, R. (2020). Evaluating contributions of recent tracking-based animal movement ecology to conservation management. *Frontiers in Ecology and Evolution*, 7, 519.
- Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye on life and planet. *Science*, 348(6240), aaa2478.
- Keller, I., Wagner, C. E., Greuter, L., Mwaiko, S., Selz, O. M., Sivasundar, A., Wittwer, S., & Seehausen, O. (2013). Population genomic signatures of divergent adaptation, gene flow and hybrid speciation in the rapid radiation of Lake Victoria cichlid fishes. *Molecular Ecology*, 22(11), 2848–2863.
- Kelly, A. C., Mateus-Pinilla, N. E., Brown, W., Ruiz, M. O., Douglas, M. R., Douglas, M. E., Shelton, P., Beissel, T., & Novakofski, J. (2014). Genetic assessment of environmental features that influence deer dispersal: Implications for prion-infected populations. *Population Ecology*, 56(2), 327–340.
- Kimmig, S. E., Beninde, J., Brandt, M., Schleimer, A., Kramer-Schadt, S., Hofer, H., Börner, K., Schulze, C., Wittstatt, U., Heddergott, M., Halczok, T., Staubach, C., & Frantz, A. C. (2020). Beyond the landscape: Resistance modelling infers physical and behavioural gene flow barriers to a mobile carnivore across a metropolitan area. *Molecular Ecology*, 29(3), 466–484.
- Knoche, S., & Lupi, F. (2012). The economic value of publicly accessible deer hunting land. *Journal of Wildlife Management*, 76(3), 462–470.
- Kopelman, N. M., Mayzel, J., Jakobsson, M., Rosenberg, N. A., & Mayrose, I. (2015). Clumpak: A program for identifying clustering modes and packaging population structure inferences across K. *Molecular Ecology Resources*, 15(5), 1179–1191.
- Kozakiewicz, C. P., Burridge, C. P., Funk, W. C., VandeWoude, S., Craft, M. E., Crooks, K. R., Ernest, H. B., Fountain-Jones, N. M., & Carver, S. (2018). Pathogens in space: Advancing understanding of pathogen dynamics and disease ecology through landscape genetics. *Evolutionary Applications*, 11(10), 1763–1778.
- Laikre, L., Schwartz, M. K., Waples, R. S., & Ryman, N. (2010). Compromising genetic diversity in the wild: Unmonitored large-scale release of plants and animals. *Trends in Ecology and Evolution*, 25(9), 520–529.
- Landguth, E. L., Cushman, S. A., Schwartz, M. K., McKelvey, K. S., Murphy, M., & Luikart, G. (2010). Quantifying the lag time to detect barriers in landscape genetics. *Molecular Ecology*, 19(19), 4179–4191.
- Lawson, D. J., van Dorp, L., & Falush, D. (2018). A tutorial on how not to over-interpret STRUCTURE and ADMIXTURE bar plots. *Nature Communications*, 9(1), 3258.
- Leberg, P. L., & Ellsworth, D. L. (1999). Further evaluation of the genetic consequences of translocations on southeastern white-tailed deer populations. *The Journal of Wildlife Management*, 63(1), 327–334.
- Leberg, P. L., Stangel, P. W., Hillestad, H. O., Marchinton, R. L., & Smith, M. H. (1994). Genetic structure of reintroduced wild turkey and white-tailed deer populations. *The Journal of Wildlife Management*, 58(4), 698.
- Leiss, W., Westphal, M., Tyshenko, M. G., Croteau, M. C., Oraby, T., Adamowicz, W., Goddard, E., Cashman, N. R., Darshan, S., & Krewski, D. (2017). Challenges in managing the risks of chronic wasting disease. *International Journal of Global Environmental Issues*, 16(4), 277–302.
- Lemopoulos, A., Prokkola, J. M., Uusi-Heikkilä, S., Vasemägi, A., Huusko, A., Hyvärinen, P., Koljonen, M.-L., Koskineniemi, J., & Vainikka, A. (2019). Comparing RADseq and microsatellites for estimating genetic diversity and relatedness – Implications for brown trout conservation. *Ecology and Evolution*, 9(4), 2106–2120.
- Leopold, A. (1931). Report on a game survey of the north central states, Sporting Arms and Ammunition Manufacturers' Institute.
- Linck, E., & Battey, C. J. (2019). Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. *Molecular Ecology Resources*, 19(3), 639–647.
- Linck, E., Epperly, K., Van Els, P., Spellman, G. M., Bryson, R. W., McCormack, J. E., Canales-Del-Castillo, R., & Klicka, J. (2019). Dense geographic and genomic sampling reveals paraphyly and a cryptic lineage in a classic sibling species complex. *Systematic Biology*, 68(6), 956–966.
- Locher, A., Scribner, K. T., Moore, J. A., Murphy, B., & Kanefsky, J. (2015). Influence of landscape features on spatial genetic structure of white-tailed deer in human-altered landscapes. *Journal of Wildlife Management*, 79(2), 180–194.
- Long, E. S., Diefenbach, D. R., Rosenberry, C. S., & Wallingford, B. D. (2008). Multiple proximate and ultimate causes of natal dispersal in white-tailed deer. *Behavioral Ecology*, 19(6), 1235–1242.
- Lopez, J., Hurwood, D., Dryden, B., & Fuller, S. (2014). Feral pig populations are structured at fine spatial scales in tropical Queensland, Australia. *PLoS One*, 9(3).
- Lowe, W. H., & Allendorf, F. W. (2010). What can genetics tell us about population connectivity? *Molecular Ecology*, 19(15), 3038–3051.
- Lundgren, E., & Ralph, P. L. (2019). Are populations like a circuit? Comparing isolation by resistance to a new coalescent-based method. *Molecular Ecology Resources*, 19(6), 1388–1406.
- Lutz, C. L., Diefenbach, D. R., & Rosenberry, C. S. (2015). Population density influences dispersal in female white-tailed deer. *Journal of Mammalogy*, 96(3), 494–501.
- Martin, B. T., Douglas, M. R., Chafin, T. K., Placyk, J. S., Birkhead, R. D., Phillips, C. A., & Douglas, M. E. (2020). Contrasting signatures of introgression in North American box turtle (*Terrapene* spp.) contact zones. *Molecular Ecology*, 29(21), 4186–4202.

- Mawdsley, J. R. (2020). Phylogenetic patterns suggest broad susceptibility to chronic wasting disease across Cervidae. *Wildlife Society Bulletin*, 44(1), 152–155.
- Mazet, O., Rodríguez, W., Grusea, S., Boitard, S., & Chikhi, L. (2016). On the importance of being structured: Instantaneous coalescence rates and human evolution-lessons for ancestral population size inference? *Heredity*, 116(4), 362–371.
- McCoy, J. E., Hewitt, D. G., & Bryant, F. C. (2005). Dispersal by yearling male white-tailed deer and implications for management. *Journal of Wildlife Management*, 69(1), 366–376.
- Meeker, R., Brown, J., Carner, B., Stephens, K., Dugger, G., Groves, B., & White, D. J. (2019). *Arkansas game and fish strategic deer management plan*. Arkansas Game and Fish Commission.
- Meffe, G. K., & Vrijenhoek, R. C. (1988). Conservation genetics in the management of desert fishes. *Conservation Biology*, 2(2), 157–169.
- Meirmans, P. G. (2012). The trouble with isolation by distance. *Molecular Ecology*, 21(12), 2839–2846.
- Miller, W. L., Miller-Butterworth, C. M., Diefenbach, D. R., & Walter, W. D. (2020). Assessment of spatial genetic structure to identify populations at risk for infection of an emerging epizootic disease. *Ecology and Evolution*, 10(9), 3977–3990.
- Miller, M. W., Runge, J. P., Holland, A. A., & Eckert, M. D. (2020). Hunting pressure modulates prion infection risk in mule deer herds. *Journal of Wildlife Diseases*, 56(4), 781–790.
- Mitchell, C. J., & Smith, W. P. (1991). Reliability of techniques for determining age in southern white-tailed deer. *Journal of the Tennessee Academy of Science*, 66(3), 117–120.
- Müller-Schwarze, D., & Müller-Schwarze, C. (1975). Subspecies specificity of response to a mammalian social odor. *Journal of Chemical Ecology*, 1(1), 125–131.
- Musmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2020). AdmixPipe: Population analyses in ADMIXTURE for non-model organisms. *BMC Bioinformatics*, 21, 337.
- Nagylaki, T. (1976). Clines with variable migration. *Genetics*, 83(4), 867–886.
- Nei, M., Maruyama, T., & Chakraborty, R. (2014). The bottleneck effect and genetic variability in populations. *Evolution*, 29(1), 1–10.
- Nelson, M. E. (1993). Natal dispersal and gene flow in white-tailed deer in northeastern Minnesota. *Journal of Mammalogy*, 74(2), 316–322.
- Nixon, C. M., Hansen, L. P., Brewer, P. A., & Chelsvig, J. E. (1991). Ecology of white-tailed deer in an intensively farmed region of Illinois. *Wildlife Monographs*, 118, 3–77.
- Nixon, C. M., & Mankin, P. C. (2016). Social and environmental influences on female white-tailed deer dispersal behavior. *Great Plains Research*, 26(2), 85–92.
- Pavlacky, D. C., Goldizen, A. W., Prentis, P. J., Nicholls, J. A., & Lowe, A. J. (2009). A landscape genetics approach for quantifying the relative influence of historic and contemporary habitat heterogeneity on the genetic connectivity of a rainforest bird. *Molecular Ecology*, 18(14), 2945–2960.
- Peischl, S., Dupanloup, I., Bosshard, L., & Excoffier, L. (2016). Genetic surfing in human populations: From genes to genomes. *Current Opinion in Genetics and Development*, 41, 53–61.
- Peterson, B. E., Storm, D. J., Norton, A. S., & Van Deelen, T. R. (2017). Landscape influence on dispersal of yearling male white-tailed deer. *Journal of Wildlife Management*, 81(8), 1449–1456.
- Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One*, 7(5), e37135.
- Petkova, D., Novembre, J., & Stephens, M. (2015). Visualizing spatial population structure with estimated effective migration surfaces. *Nature Genetics*, 48(1), 94–100.
- Polechová, J., & Barton, N. (2011). Genetic drift widens the expected cline but narrows the expected cline width. *Genetics*, 189(1), 227–235.
- Potapov, A., Merrill, E., Pybus, M., & Lewis, M. A. (2016). Chronic wasting disease: Transmission mechanisms and the possibility of harvest management. *PLoS One*, 11(3), e0151039.
- Puckett, E. E. (2017). Variability in total project and per sample genotyping costs under varying study designs including with microsatellites or SNPs to answer conservation genetic questions. *Conservation Genetics Resources*, 9(2), 289–304.
- Puckett, E. E., & Eggert, L. S. (2016). Comparison of SNP and microsatellite genotyping panels for spatial assignment of individuals to natal range: A case study using the American black bear (*Ursus americanus*). *Biological Conservation*, 193, 86–93.
- Rengefors, K., Gollnisch, R., Sassenhagen, I., Härnström Aloisi, K., Svensson, M., Leuret, K., Čertnerová, D., Cresko, W. A., Bassham, S., & Ahrén, D. (2021). Genome-wide single nucleotide polymorphism markers reveal population structure and dispersal direction of an expanding nuisance algal bloom species. *Molecular Ecology*, 30(4), 912–925.
- Richardson, J. L., Brady, S. P., Wang, I. J., & Spear, S. F. (2016). Navigating the pitfalls and promise of landscape genetics. *Molecular Ecology*, 25, 849–863.
- Rivera, N. A., Brandt, A. L., Novakofski, J. E., & Mateus-Pinilla, N. E. (2019). Chronic wasting disease in cervids: Prevalence, impact and management strategies. *Veterinary Medicine: Research and Reports*, 10, 123–139.
- Roberts, G. O., Gelman, A., & Gilks, W. R. (1997). Weak convergence and optimal scaling of random walk Metropolis algorithms. *Annals of Applied Probability*, 7(1), 110–120.
- Robinson, S. J., Samuel, M. D., Lopez, D. L., & Shelton, P. (2012). The walk is never random: Subtle landscape effects shape gene flow in a continuous white-tailed deer population in the Midwestern United States. *Molecular Ecology*, 21(17), 4190–4205.
- Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: A versatile open source tool for metagenomics. *PeerJ*, 4, e2409v1.
- Rousset, F. (1997). Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics*, 145, 1219–1228.
- Rubin, B. E. R., Ree, R. H., & Moreau, C. S. (2012). Inferring phylogenies from RAD sequence data. *PLoS One*, 7(4), e33394.
- Sacks, B. N., Brazeal, J. L., & Lewis, J. C. (2016). Landscape genetics of the nonnative red fox of California. *Ecology and Evolution*, 6(14), 4775–4791.
- Santangelo, J. S., Johnson, M. T. J., & Ness, R. W. (2018). Modern spandrels: The roles of genetic drift, gene flow and natural selection in the evolution of parallel clines. *Proceedings of the Royal Society B: Biological Sciences*, 285(1878), 20180230.
- Schoville, S. D., Bonin, A., François, O., Lobreaux, S., Melodelima, C., & Manel, S. (2012). Adaptive genetic variation on the landscape: Methods and cases. *Annual Review of Ecology, Evolution, and Systematics*, 43(1), 23–43.
- Seddon, P. J., Griffiths, C. J., Soorae, P. S., & Armstrong, D. P. (2014). Reversing defaunation: Restoring species in a changing world. *Science*, 345(6195), 406–412.
- Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., Pagad, S., Pyšek, P., Winter, M., Arianoutsou, M., Bacher, S., Blasius, B., Brundu, G., Capinha, C., Celesti-Grappo, L., Dawson, W., Dullinger, S., Fuentes, N., Jäger, H., ... Essl, F. (2017). No saturation in the accumulation of alien species worldwide. *Nature Communications*, 8, 1–9.
- Severinghaus, C. W. (1949). Tooth development and wear as criteria of age in white-tailed deer. *The Journal of Wildlife Management*, 13, 195–216.
- Shaw, J. C., Lancia, R. A., Conner, M. S., & Rosenberry, C. S. (2006). Effect of population demographics and social pressures on white-tailed deer dispersal ecology. *Journal of Wildlife Management*, 70(5), 1293–1301.
- Shephard, J. M., Ogden, R., Tryjanowski, P., Olsson, O., & Galbusera, P. (2013). Is population structure in the European white stork

- determined by flyway permeability rather than translocation history? *Ecology and Evolution*, 3(15), 4881–4895.
- Short Bull, R. A., Cushman, S. A., Mace, R., Chilton, T., Kendall, K. C., Landguth, E. L., Schwartz, M. K., Mckelvey, K., Allendorf, F. W., & Luikart, G. (2011). Why replication is important in landscape genetics: American black bear in the Rocky Mountains. *Molecular Ecology*, 20(6), 1092–1107.
- Simpson, E. H. (1949). Measurement of diversity. *Nature*, 163(1943), 688.
- Slatkin, M. (1973). Gene flow and selection in a cline. *Genetics*, 75(4), 733–756.
- Sumners, J. A., Demarais, S., DeYoung, R. W., Honeycutt, R. L., Rooney, A. P., Gonzales, R. A., & Gee, K. L. (2015). Variable breeding dates among populations of white-tailed deer in the southern United States: The legacy of restocking? *Journal of Wildlife Management*, 79(8), 1213–1225.
- Travis, J. M. J., Münkemüller, T., Burton, O. J., Best, A., Dytham, C., & Johst, K. (2007). Deleterious mutations can surf to high densities on the wave front of an expanding population. *Molecular Biology and Evolution*, 24(10), 2334–2343.
- Uehlinger, F. D., Johnston, A. C., Bollinger, T. K., & Waldner, C. L. (2016). Systematic review of management strategies to control chronic wasting disease in wild deer populations in North America. *BMC Veterinary Research*, 12(1), 173.
- Vasemägi, A. (2006). The adaptive hypothesis of clinal variation revisited: Single-locus clines as a result of spatially restricted gene flow. *Genetics*, 173(4), 2411–2414.
- Waller, D. M., & Alverson, W. S. (1997). The white-tailed deer: A keystone herbivore. *Wildlife Society Bulletin*, 25(2), 217–226.
- Williams, E. S., & Young, S. (1980). Chronic wasting disease of captive mule deer: A spongiform encephalopathy. *Journal of Wildlife Diseases*, 16(1), 89–98.
- Wolverton, S., Kennedy, J. H., & Cornelius, J. D. (2007). A paleozoological perspective on white-tailed deer (*Odocoileus virginianus texana*) population density and body size in central Texas. *Environmental Management*, 39(4), 545–552.
- Wood, R. (1944). Arkansas' deer transplanting program. *Transactions of the North American Wildlife Conference*, 9, 162–167.
- Wynn, S. (1943). Howard county game refuge and deer farm. *The Arkansas Historical Quarterly*, 2(4), 340–345.
- Zalewski, A., Piertney, S. B., Zalewska, H., & Lambin, X. (2009). Landscape barriers reduce gene flow in an invasive carnivore: Geographical and local genetic structure of American mink in Scotland. *Molecular Ecology*, 18(8), 1601–1615.
- Zellmer, A. J., & Knowles, L. L. (2009). Disentangling the effects of historic vs. contemporary landscape structure on population genetic divergence. *Molecular Ecology*, 18(17), 3593–3602.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Chafin TK, Zbinden ZD, Douglas MR, et al. Spatial population genetics in heavily managed species: Separating patterns of historical translocation from contemporary gene flow in white-tailed deer. *Evol Appl.* 2020;00:1–17. <https://doi.org/10.1111/eva.13233>